

# Search for Gauge-Mediated Supersymmetry-Breaking in Diphoton Events with Missing Transverse Energy at CDF II

(Dated: July 24, 2009; Version 0.5.3; Line-count 464 out of 460)

We present the results of search for gauge-mediated supersymmetry-breaking with  $\tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}$  in the  $\gamma\gamma$ +missing transverse energy final state. In  $2.6 \pm 0.2 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s}=1.96 \text{ TeV}$  recorded by the CDF II detector we observe no candidate events, consistent with a standard model background expectation of  $1.4 \pm 0.4$  events. We set 95% C.L. cross section limits and place the world-best limit on the  $\tilde{\chi}_1^0$  mass of  $149 \text{ GeV}/c^2$  at  $\tau_{\tilde{\chi}_1^0}=0 \text{ ns}$  as well as make exclusions in the  $\tilde{\chi}_1^0$  mass-lifetime plane for  $\tau_{\tilde{\chi}_1^0} \lesssim 2 \text{ ns}$ .

PACS numbers: 13.85.Rm, 12.60.Jv, 13.85.Qk, 14.80.Ly

The standard model (SM) of elementary particles has been enormously successful, but is incomplete. Theoretical motivation [1] and the observation of the ‘ $ee\gamma\gamma$ +missing transverse energy ( $\cancel{E}_T$ )’ [2, 3] candidate event by the CDF experiment during Run I at the Fermilab Tevatron provide compelling rationale to search for the production and decay of new heavy particles that produce events with final state photons and  $\cancel{E}_T$  in collider experiments. Of particular theoretical interest are supersymmetry (SUSY) models with gauge-mediated SUSY-breaking (GMSB) [1]. These models solve the “naturalness problem” [4] and provide a low mass (warm) dark matter candidate that is both consistent with inflation and astronomical observations [5]. Since many versions of these models have a similar phenomenology, we consider the scenario in which the lightest neutralino ( $\tilde{\chi}_1^0$ ) decays almost exclusively into a photon ( $\gamma$ ) and a weakly interacting, stable gravitino ( $\tilde{G}$ ) that gives rise to  $\cancel{E}_T$  by leaving a detector without depositing any energy [6]. In these models, the  $\tilde{\chi}_1^0$  is favored to have a lifetime on the order of a nanosecond, and the  $\tilde{G}$  is a warm dark matter candidate with a mass in the range  $0.5 \text{ keV} < m_{\tilde{G}} < 1.5 \text{ keV}$  [7]. Other direct searches [8–10] have constrained the mass of the  $\tilde{\chi}_1^0$  to  $\gtrsim 100 \text{ GeV}/c^2$  for much of the parameter space. At the Tevatron, sparticle production is dominated by gaugino pairs, and the  $\tilde{\chi}_1^0$  mass ( $m_{\tilde{\chi}_1^0}$ ) and lifetime ( $\tau_{\tilde{\chi}_1^0}$ ) are the two most important parameters in determining the final states and their kinematics. Different search strategies are required for  $\tilde{\chi}_1^0$  lifetimes above and below about a nanosecond [11].

In this letter we describe a search for GMSB with  $\tau_{\tilde{\chi}_1^0} \leq 2 \text{ ns}$ , favored for large  $m_{\tilde{\chi}_1^0}$ , in which gaugino pairs are produced and decay to the  $\gamma\gamma + \cancel{E}_T + X$  final state where  $X$  denote other high  $E_T$  final state particles. We use a dataset corresponding to an integrated luminosity of  $2.6 \pm 0.2 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s}=1.96 \text{ TeV}$  from the Tevatron collected with the CDF II detector [12]. This is ten times more data than our previous search [8]. For the first time in this channel we use our new photon timing system [13] and a new model of the  $\cancel{E}_T$  resolution (METMODEL) [14]. These additions significantly enhance our rejection of backgrounds from instrumental and non-collision sources, which allows our search to considerably extend the sensitivity for large  $\tilde{\chi}_1^0$  mass compared to other Tevatron searches [9]. It

is the first to consider the  $\tau_{\tilde{\chi}_1^0}$  up to 2 ns.

Our strategy is to select  $\gamma\gamma$  candidates and optimize the search for the presence of both significant  $\cancel{E}_T$  and large total event transverse energy to indicate the decays of heavy gauginos. We perform an *a priori* analysis based on the expected sensitivity, taking into account the background and signal predictions.

Here we briefly describe the aspects of the detector [12] relevant to this analysis. The magnetic spectrometer consists of tracking devices inside a 3-m diameter, 5-m long superconducting solenoid magnet that operates at 1.4 T. A 3.1-m long drift chamber (COT) with 96 layers of sense wires measures the  $z$  position and time of the  $p\bar{p}$  interaction and the momenta of charged particles. The calorimeter consists of projective towers with electromagnetic (EM) and hadronic (HAD) compartments and is divided into a central barrel that surrounds the solenoid coil ( $|\eta| < 1.1$ ) [2] and a pair of end-plugs that cover the region  $1.1 < |\eta| < 3.6$ . Both are used to identify and measure the 4-momenta of photons, electrons, jets ( $j$ ) [15] and provide  $\cancel{E}_T$  information. The EM calorimeter is instrumented with a timing system, EMTiming [13], that measures the arrival time of photons.

Our analysis begins with events passing the CDF three-level trigger. The combined trigger selection efficiency is effectively 100% for our diphoton events [14]. We then select events with two photons with  $|\eta| < 1.1$  and  $E_T > 13$  GeV. Offline, both photons are required to be in the fiducial part of the calorimeter and pass the standard CDF photon identification (ID) and isolation requirements [8] with two minor modifications to remove instrumental and electron backgrounds [14, 16]. These events are dominated by SM production of  $\gamma\gamma$ ,  $\gamma j$  with  $j \rightarrow \gamma_{fake}$  and  $jj \rightarrow \gamma_{fake}\gamma_{fake}$ , where  $\gamma_{fake}$  is a jet misidentified as a photon. To minimize the number of these events with large  $\cancel{E}_T$  due to calorimeter energy mismeasurements we remove events if the  $\cancel{E}_T$  vector points within  $|\Delta\phi| < 0.3$  of the second highest  $E_T$  photon or any jet pointing to the calorimeter cracks [14]. To help maintain the projective nature of the photon reconstruction in the calorimeter we require a vertex with  $|z_{vertex}| < 60$  cm, which also helps to reduce non-collision backgrounds. For events with multiple reconstructed vertices we pick the vertex with the highest  $\sum_{tracks} p_T$  [2], unless assigning the photons to a different vertex lowers the  $\cancel{E}_T$ , in which case we choose that vertex.

Non-collision backgrounds coming from cosmic rays and beam-related effects can produce  $\gamma\gamma + \cancel{E}_T$  candidates, and are removed from the inclusive  $\gamma\gamma$  sample using a number of techniques. Photon candidates from cosmic rays are not correlated in time with collisions and events are removed if the timing of either photon, corrected for average path length ( $t_\gamma$ ), indicates a non-collision source being from the collision [14, 16]. Photon candidates can also be produced by beam-related muons that originate upstream of the detector. These are suppressed using standard beam halo identification requirements [16]. A total of 38,053 inclusive  $\gamma\gamma$  candidate events pass our requirements.

Backgrounds to the  $\gamma\gamma + \cancel{E}_T$  final state from SM  $\gamma\gamma / \gamma\gamma_{fake} / \gamma_{fake}\gamma_{fake}$  and mismeasured (“fake”)  $\cancel{E}_T$  arise due to energy mismeasurements in the calorimeter or event reconstruction pathologies. To select events with real and significant  $\cancel{E}_T$ , as part of the optimization, and to predict the contribution of SM backgrounds with fake  $\cancel{E}_T$  due to normal energy measurement fluctuations, we use METMODEL [14]. This algorithm considers the clustered (jets) and unclustered energy in the event and calculates a probability for fluctuations in the energy measurement to produce

1  $\cancel{E}_T^{fluct}$  equivalent to or larger than the measured  $\cancel{E}_T$  ( $P_{\cancel{E}_T^{fluct} \geq \cancel{E}_T}$ ). This probability is then used to define a  $\cancel{E}_T$ -  
 2 significance as  $-\log_{10} \left( P_{\cancel{E}_T^{fluct} \geq \cancel{E}_T} \right)$ . Events with true and fake  $\cancel{E}_T$  of the same value have, on average, different  
 3  $\cancel{E}_T$ -significance. To estimate the expected  $\cancel{E}_T$ -significance distribution for SM events with fake  $\cancel{E}_T$ , and the number  
 4 of mismeasured events above a given  $\cancel{E}_T$ -significance requirement, we use pseudo-experiments in which we smear the  
 5 jets and unclustered energy according to their resolution functions in the event. The systematic uncertainty in the  
 6 METMODEL is dominated by the uncertainty in the resolution functions [14].

7 Reconstruction pathologies in SM events with no intrinsic  $\cancel{E}_T$ , such as a wrong choice of the primary interaction  
 8 vertex or tri-photon events with a lost photon, are unaccounted for by the METMODEL. To obtain the prediction for  
 9 this background we model SM kinematics and event reconstruction using a PYTHIA [17]  $\gamma\gamma$  sample with a detector  
 10 simulation [18]. This sample is normalized to the number of events in the inclusive  $\gamma\gamma$  data sample to take into account  
 11  $\gamma j$  and  $jj$  contributions to the backgrounds. We subtract the expectations for energy mismeasurement fluctuations  
 12 in the MC to avoid double counting. Uncertainties are dominated by the statistics of the MC sample.

13 Electroweak production of  $W$ 's and  $Z$ 's which decay to leptons can also give rise to the  $\gamma\gamma + \cancel{E}_T$  signature where  
 14 one or more of the photons can be fake, but the  $\cancel{E}_T$  is due to one or more neutrinos. To estimate the contribution  
 15 from these backgrounds we use MC simulations, normalized to their production cross sections and considering all  
 16 the leptonic decay modes of the bosons. The Baur MC [19] is used to simulate  $W\gamma$  and  $Z\gamma$  production and decay  
 17 where initial and final state radiation (ISR/FSR) simulates  $W/Z + \gamma\gamma$  events. The PYTHIA [17] MC is used to simulate  
 18 backgrounds where both photons are fakes:  $W$ ,  $Z$ , with photons from ISR/FSR removed, and  $t\bar{t}$  sources. To minimize  
 19 the dependence of our predictions on potential MC-Data differences we scale our MC predictions to the observed  
 20 number of  $e\gamma$  events [14] in data where we use the same diphoton triggers and analysis selection procedures developed  
 21 for our inclusive  $\gamma\gamma$  sample. Uncertainties are dominated by the statistics of the MC and  $e\gamma$  normalization data  
 22 sample.

23 Non-collision backgrounds are estimated using the data. Using the sample selection requirements, but requiring  
 24 one of the photons to have  $t_\gamma > 25$  ns we identify a cosmic enhanced sample. Similarly, we select a beam related  
 25 background enhanced sample. We estimate the number of these events in the signal region using the ratio of events  
 26 outside the timing requirements to events inside the signal region and the measured efficiencies of the non-collision  
 27 rejection requirements [14]. The uncertainties on both non-collision background estimates are dominated by the  
 28 statistical uncertainty on the number of identified events. The top of Figure 1 shows the  $\cancel{E}_T$ -significance distribution  
 29 for the inclusive  $\gamma\gamma$  sample, along with the predictions for all the backgrounds.

30 We estimate the sensitivity to heavy, neutral particles that decay to photons using the GMSB reference model [6]  
 31 in the mass-lifetime range,  $75 \leq m_{\tilde{\chi}_1^0} \leq 150$  GeV and  $\tau_{\tilde{\chi}_1^0} \leq 2$  ns. Events from all SUSY processes [16] are simulated  
 32 with PYTHIA [17] followed by a detector simulation [18]. The acceptance is a strong function of the fraction of  $\tilde{\chi}_1^0$   
 33 decays that occur in the detector volume, which is dependent on both the  $\tau_{\tilde{\chi}_1^0}$  and the masses of the sparticles, all




FIG. 1: The top plot shows the  $E_T$ -significance distribution for the inclusive  $\gamma\gamma$  candidate sample, along with the background predictions. The bottom plot shows the predicted  $H_T$  (total  $E_T$  of photons, jets and  $E_T$ ) distribution after all but the final  $H_T$  requirement.

of which scale linearly with  $m_{\tilde{\chi}_1^0}$  for our model [11]. The total systematic uncertainty on the acceptance, after all kinematic requirements, is estimated to be 7%, dominated by the uncertainty in the photon ID efficiency (2.5% per photon). Other significant contributions come from uncertainties on ISR/FSR (4%), jet energy measurement (2%),  $E_T$ -significance parameterizations (1%) and the parton distribution functions (1%).

We determine the final kinematic selection requirements by optimizing the mean expected 95% confidence level (C.L.) cross section limit in the no-signal assumption without looking at the data in the signal region [20]. To compute the predicted cross section upper limit we combine the luminosity, the acceptance and the background

estimates with their systematic uncertainties using a Bayesian method [21]. The predicted limits are optimized by simultaneously varying the selection requirements for  $\cancel{E}_T$ -significance,  $H_T$  (scalar sum of  $E_T$  of photons, jets and  $\cancel{E}_T$ ), and the azimuthal angle between the two leading photons,  $\Delta\phi(\gamma_1, \gamma_2)$ . The large  $\cancel{E}_T$ -significance requirement eliminates most of the SM background with fake  $\cancel{E}_T$ . GMSB production is dominated by heavy gaugino pairs which decay to high  $E_T$ , light final state particles via cascade decays. GMSB signal has, on average, larger  $H_T$  compared to SM backgrounds so that an  $H_T$  requirement can remove these backgrounds effectively. Electroweak backgrounds with large  $H_T$  typically consist of a high  $E_T$  photon recoiling against  $W \rightarrow e\nu$ , identified as  $\gamma_{fake}\cancel{E}_T$ , which means the gauge boson decay is highly boosted. Thus, the two photon candidates in the final state are mostly back-to-back. Also, the high  $E_T$  diphotons with large  $H_T$  from SM background are mostly back-to-back with fake  $\cancel{E}_T$ ; the  $\Delta\phi(\gamma_1, \gamma_2)$  cut, therefore, reduces both these backgrounds.

While each point in the considered  $\tau_{\tilde{\chi}_1^0}$  vs.  $m_{\tilde{\chi}_1^0}$  space gives a slightly different optimization, we choose a single set of requirements to maximize the expected 95% C.L. exclusion region; where the predicted production cross section at next-to-leading order [22] is above the expected cross section limit. The exclusion region also takes into account the production cross section uncertainties which are dominated by the parton distribution functions (7%) and the renormalization scale (3%). We find the optimal set of cuts, before unblinding the signal region, to be:  $\cancel{E}_T\text{-significance} > 3$ ,  $H_T > 200$  GeV, and  $\Delta\phi(\gamma_1, \gamma_2) < \pi - 0.35$ . With these requirements we predict  $1.4 \pm 0.4$  background events with  $0.9 \pm 0.4$  events from electroweak sources (dominated by  $Z\gamma$  production) with real  $\cancel{E}_T$ ,  $0.5 \pm 0.2$  from SM with fake  $\cancel{E}_T$ , and  $0.001^{+0.008}_{-0.001}$  from non-collision sources. The acceptance for  $m_{\tilde{\chi}_1^0} = 140$  GeV/c<sup>2</sup> and  $\tau_{\tilde{\chi}_1^0} = 0$  ns is estimated to be  $7.8 \pm 0.6\%$ .

No events in the data pass the final event selection. The predicted  $H_T$  distribution is shown in Figure 1 (bottom), after all but the final  $H_T$  cut. The data is consistent with the no-signal hypothesis and well modeled by backgrounds alone. We set cross section limits as a function of  $m_{\tilde{\chi}_1^0}$  and  $\tau_{\tilde{\chi}_1^0}$  respectively, as shown in Figure 2. The  $m_{\tilde{\chi}_1^0}$  reach, based on the predicted and observed number of events for  $\tau_{\tilde{\chi}_1^0} = 0$ , is 141 GeV/c<sup>2</sup> and 149 GeV/c<sup>2</sup> respectively. These limits significantly extend the search sensitivity beyond the results of D0 [9] for  $\tau_{\tilde{\chi}_1^0} \leq 2$  ns and, when combined with the complementary exclusions from CDF and LEP [10, 16], cover the region shown in Figure 3.

In conclusion, we have performed an optimized search for heavy, neutral particles that decay to photons in the  $\gamma\gamma + \cancel{E}_T$  final state using 2.6 fb<sup>-1</sup> of data. There is no excess of events beyond expectations. We set cross section limits using a GMSB model with  $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$ , and find an exclusion region in the  $\tau_{\tilde{\chi}_1^0}$ - $m_{\tilde{\chi}_1^0}$  plane with a world-best 95% C.L. lower limit on the  $\tilde{\chi}_1^0$  mass of 149 GeV/c<sup>2</sup> at  $\tau_{\tilde{\chi}_1^0} = 0$  ns. By the end of Run II, an integrated luminosity of 10 fb<sup>-1</sup> is possible for which we estimate a mass reach of  $\simeq 160$  GeV/c<sup>2</sup> at a lifetime of 0 ns.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the




FIG. 2: The predicted and observed 95% C.L. cross section upper limits as a function of the  $\tilde{\chi}_1^0$  mass at a lifetime of 0 ns (top) and as a function of the  $\tilde{\chi}_1^0$  lifetime at a mass of 140 GeV/ $c^2$  (bottom). Indicated in green (darker shading) is the production cross section, along with its 8.0% uncertainty-band. In yellow (lighter shading) is the RMS variation on the expected cross section limit.

- 1 Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung,
- 2 Germany; the Korean Science and Engineering Foundation and the Korean Research Foundation; the Particle Physics
- 3 and Astronomy Research Council and the Royal Society, UK; the Russian Foundation for Basic Research; the Comisión
- 4 Interministerial de Ciencia y Tecnología, Spain; in part by the European Community's Human Potential Programme
- 5 under contract HPRN-CT-2002-00292; and the Academy of Finland.




FIG. 3: The predicted and observed exclusion region along with the limit from ALEPH/LEP [10] and the CDF  $\gamma + \cancel{E}_T + \text{jet}$  ‘delayed’ photon analysis [16]. We have a mass reach of  $141 \text{ GeV}/c^2$  (predicted) and  $149 \text{ GeV}/c^2$  (observed) at lifetimes up to  $1 \text{ ns}$ . The shaded band shows the parameter space where  $0.5 < m_{\tilde{G}} < 1.5 \text{ keV}/c^2$ , favored by cosmological models [7].

- 
- [1] S. Dimopoulos, S. Thomas, J.D. Wells, Nucl. Phys. B **488**, 39 (1997); S. Ambrosanio, G.D. Kribs and S.P. Martin, Phys. Rev. D **56**, 1761 (1997); G.F. Giudice and R. Rattazzi, Phys. Rept. **322**, 419 (1999); and S. Ambrosanio, G. Kane, G. Kribs, S. Martin and S. Mrenna, Phys. Rev. D **55**, 1372 (1997).
- [2] We use a cylindrical coordinate system in which the proton beam travels along the  $z$ -axis,  $\theta$  is the polar angle,  $\phi$  is the azimuthal angle relative to the horizontal plane, and  $\eta = -\ln \tan(\theta/2)$ . The transverse energy and momentum are defined as  $E_T = E \sin \theta$  and  $p_T = p \sin \theta$  where  $E$  is the energy measured by the calorimeter and  $p$  the momentum measured in the tracking system.  $\cancel{E}_T = |-\sum_i E_T^i \vec{n}_i|$  where  $\vec{n}_i$  is a unit vector that points from the interaction vertex to the  $i^{\text{th}}$  calorimeter

tower in the transverse plane.

- [3] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **81**, 1791 (1998) and Phys. Rev. D **59**, 092002 (1999).
- [4] S. Martin, arXiv:hep-ph/9709356.
- [5] P. Bode, J. Ostriker, and N. Turok, Astrophys. J. **556**, 93 (2001).
- [6] B.C. Allanach *et al.*, Eur. Phys. J. C **25**, 113 (2002). We use benchmark model 8 and take the messenger mass scale  $M_{\text{mess}}=2\Lambda$ ,  $\tan(\beta)=15$ ,  $\mu>0$  and the number of messenger fields  $N_{\text{mess}}=1$ . The  $\tilde{G}$  mass factor and the supersymmetry breaking scale  $\Lambda$  are allowed to vary independently.
- [7] C.-H. Chen and J.F. Gunion, Phys. Rev. D **58**, 075005 (1998).
- [8] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 031104 (2005).
- [9] V.M. Abazov *et al.* (D0 Collaboration), Phys. Lett. B **659**, 856 (2008).
- [10] A. Heister *et al.* (ALEPH Collaboration), Eur. Phys. J. C **25**, 339 (2002); also see M. Gataullin, S. Rosier, L. Xia and H. Yang, arXiv:hep-ex/0611010; G. Abbiendi *et al.* (OPAL Collaboration), Proc. Sci. HEP2005 346 (2006); J. Abdallah *et al.* (DELPHI Collaboration), Eur. Phys. J. C **38** 395 (2005).
- [11] D. Toback and P. Wagner, Phys. Rev. D **70**, 114032 (2004).
- [12] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005).
- [13] M. Goncharov *et al.*, Nucl. Instrum. Methods A **565**, 543 (2006).
- [14] T. Aaltonen *et al.* (CDF Collaboration),  $\gamma\gamma+X$  Phys. Rev. D, to be submitted. Photons with second highest  $E_T$  or narrow jets that occur in few towers and have few tracks, can be mismeasured if they are located close to the calorimeter cracks at  $\eta \sim 0$  and  $|\eta| \sim 1.1$ .
- [15] For a discussion of the jet energy measurements, see T. Affolder *et al.* (CDF Collaboration), Phys. Rev. D. **64**, 032001 (2001). For a discussion of standard jet correction systematics, see A. Bhatti *et al.*, Nucl. Instrum. Methods, A 566, 375 (2006). We use jets with cone size  $\Delta R=0.4$ .
- [16] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. Lett. **99**, 121801 (2007); T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. D **78**, 032015 (2008).
- [17] T. Sjöstrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001). We use version 6.216.
- [18] We use the standard GEANT based detector simulation [R. Brun *et al.*, CERN-DD/EE/84-1 (1987)] and add a parametrized EMTiming simulation.
- [19] U. Baur, T. Han and J. Ohnemus, Phys. Rev. D **48**, 5140 (1993); U. Baur, T. Han and J. Ohnemus, *ibid.* **57**, 2823 (1998); The  $W\gamma$  and  $Z\gamma$  processes are simulated using the leading-order event generator and have a k-factor fixed at 1.36. The initial and final state radiations (resulting in additional jets or photons), underlying event, and additional interactions are simulated by PYTHIA [17].
- [20] E. Boos, A. Vologdin, D. Toback, and J. Gaspard, Phys. Rev. D **66**, 013011 (2002).
- [21] T. Junk, Nucl. Instrum. Methods A **434**, 435-443 (1999).
- [22] We use the leading-order cross sections generated by PYTHIA [17] and the k-factors produced by PROSPINO 2.0 [W. Beenakker *et al.*, Phys. Rev. Lett. **83**, 3780 (1999)].